



# Urinary Tract Infection

## CONSULT FORM for MedAdvisor Users



*This form is intended to be used as a tool to collect information which will be entered into MedAdvisor. A service summary must be generated and provided to the patient.*

Confirm the patient is aware of charges for the service and consents to their personal information being recorded and a dispense record being uploaded to their My Health Record.

Full Name:

Date of Birth (18-65 years only):

Address:

Sex (anatomical):

Male

Female

Medicare No:

What symptoms is the patient experiencing (patient must have 2 or more of the following)?

Dysuria

Notes

Suprapubic pain

Urinary frequency

Urinary urgency

Refer if the patient has any of the following:

**YES NO**

Has any vaginal symptoms, itch, discharge?

Has any systemic symptoms (fever, chills, nausea, feeling particularly unwell, flank pain)?

Is pregnant, postpartum (4-6 weeks after birth) or had a recent miscarriage or abortion?

Has been treated for a UTI in the last 6 months?

Has been treated for more than two UTIs in the last 12 months?

Has prior non-responsiveness to UTI treatment? (e.g. symptoms of a UTI persisting 48-72 hours after starting treatment)

Had recurrence of UTI symptoms within 2 weeks of completing antimicrobial treatment?

Has been hospitalised in the last 4 weeks or been an inpatient of a health care facility in the last 3 months?

Has travelled overseas in the last 3 months or to a country with a high prevalence of antibiotic (eg. Southeast Asia and South Asia) in the last 6 months?

Is at risk of having an STI?

Is on medications that suppress the immune system (azathioprine, cyclosporin, methotrexate)?



# Urinary Tract Infection

## CONSULT FORM for MedAdvisor Users



*This form is intended to be used as a tool to collect information which will be entered into MedAdvisor. A service summary must be generated and provided to the patient.*

Refer if the patient has any of the following:

**YES NO**

- Visible haematuria (blood in the urine)
  - Pyuria (pus in the urine)
  - Urinary tract abnormality or obstruction
  - History of renal stones
  - Urinary catheter within the last 48 hours
  - Previous episodes of pyelonephritis
  - Immunocompromised
  - Spinal cord injury
  - Diabetes
  - IUD in situ
  - Antibiotic use in the past 3 months
  - History of blood disorder/porphyria
-