## Pharmacist Administered Vaccinations in the Northern Territory



This document has been developed to facilitate a quick reference for determining vaccine administration eligibility in the Northern Territory in light of recent changes to the scope of vaccinations.

Vaccines included in the National Immunisation Program can be funded through the NIPVIP program or charged privately to the patient.

Pharmacists are recommended to use this document in conjunction with the <u>National Immunisation Program schedule</u>, the <u>Australian Immunisation Handbook</u> and the <u>Pharmacist Vaccination Scheduled Substance Treatment Protocol</u>.




Figure 1: Pharmacist Administered Vaccinations in the Northern Territory Flow Chart

This flow chat has been developed to provide a quick reference for interpreting the information provided in the Pharmacist Administered Vaccinations in Table 1: Pharmacist Administered Vaccinations in the Northern Territory on Pages 2 and 3

The information in this document is current as of the publication date. This information serves as a guide only and should not be employed as a substitute or replacement for reviewing regulations or as a clinical decision-making tool. Published 13 March 2023

## Pharmacist Administered Vaccinations in the Northern Territory



**Table 1: Pharmacist Administered Vaccinations in the Northern Territory.** 

Vaccine	Pharmacists Legally able to administer in the NT	National Immunisation Program (+ Age)	NT schedule (+ Age)	NIP Eligibility Notes	Administration Notes
Influenza (flu)	<b>√</b>	<b>√</b> (5)	<b>√</b> (5)	<ul> <li>5+ years people with specified medical risk conditions</li> <li>5+ years Aboriginal and Torres Strait Islander</li> <li>65+ years non-Aboriginal and Torres Strait Islander adults</li> <li>Pregnant women</li> </ul>	All persons 5+ requesting Influenza vaccination. User pays.
Diphtheria, tetanus, pertussis (dTpa)	✓	<b>√</b> (5)	<b>√</b> (5)	<ul> <li>12–13 years</li> <li>Catch up to 20 years</li> <li>Pregnant women from 20 weeks</li> </ul>	Can be given to all persons 5+ as recommended by the Australian Immunisation Handbook. User Pays.
Measles, mumps, rubella (MMR)	✓	<b>√</b> (5)	<b>√</b> (5)	<ul> <li>Catch up 5+years up to 20 years</li> <li>Refugees aged 20yrs and over</li> </ul>	Can be given to all persons 5+ as recommended by the Australian Immunisation Handbook. User Pays.
Human papillomavirus (HPV)	✓	<b>√</b> (5)	<b>√</b> (5)	<ul><li>12–13 years</li><li>Catch up to 25 years</li></ul>	Can be given to all persons 5+ as recommended by the Australian Immunisation Handbook. User Pays.
Haemophilus influenzae type b (Hib)	<b>√</b>	<b>√</b> (5)	X	<ul> <li>5+ years with asplenia or hyposplenia,</li> <li>a single dose is required if the person was not vaccinated in infancy or incompletely vaccinated</li> <li>Catch up 5+years</li> </ul>	
Hepatitis A	<b>√</b>	<b>√</b> (5)	Х	Catch up 5+ years for Aboriginal and Torres Strait Islander people only	
Hepatitis B	<b>√</b>	<b>√</b> (5)	Х	<ul> <li>Catch up 5+ years for all Australians.</li> <li>Refugees aged 20yrs and over.</li> <li>Pharmacists require serology confirmation of inadequate coverage as per Australian Immunisation Handbook</li> </ul>	
Hepatitis A and B combination	<b>√</b>	Maybe	X	<ul> <li>Combination vaccines may be used where both Hepatitis A and B NIP criteria are met</li> </ul>	
Meningococcal disease (MenACWY)	<b>√</b>	<b>√</b> (5)	X	<ul><li>14–16 years</li><li>Catch up to 20 years</li></ul>	
Meningococcal B	✓	<b>√</b> (5)	X	<ul> <li>Catch up 5+ years</li> <li>Aboriginal and Torres Strait Islander children</li> </ul>	
Pneumococcal disease	<b>√</b>	<b>√</b> (5)	X	<ul> <li>50+ years Aboriginal and Torres Strait Islander adults</li> <li>70+ years non-Aboriginal and Torres Strait Islander adults</li> <li>Catch up 5+ years up to 20 years</li> </ul>	

## Pharmacist Administered Vaccinations in the Northern Territory



Poliomyelitis	✓	<b>√</b> (5)	X	<ul><li>Catch up 5+ years up to 20 years</li><li>Refugees aged 20yrs and over</li></ul>	
Varicella (chickenpox)	✓	<b>√</b> (5)	X	<ul><li>Catch up 5+ years up to 20 years</li><li>Refugees aged 20yrs and over</li></ul>	
Zoster (herpes zoster)	<b>√</b>	<b>√</b> (5)	x	<ul> <li>18+ years people with specified medical risk conditions</li> <li>50+ years Aboriginal and Torres Strait Islander adults</li> <li>65+ years non-Aboriginal and Torres Strait Islander adults</li> </ul>	
COVID-19 vaccines: AZ, PF, Moderna, Novavax	<b>√</b>	X	<b>√</b> (5)		Commonwealth Funded
Japanese encephalitis	✓	X	<b>√</b> (5)		Can be given to all persons 5+ as recommended by the Australian Immunisation Handbook. User Pays.
Monkeypox	✓	X	<b>√</b> (5)		Can be given to all persons 5+ as recommended by the Australian Immunisation Handbook. User Pays.
Cholera (oral)	X	X	X		
dTpa in combination with poliomyelitis (dTpa-IPV)	X	X	X		
Measles, mumps, rubella, varicella	X	Х	X		
Meningococcal C	Х	Х	Х		
Q fever	Х	Х	Х		
Rabies and other lyssaviruses (pre-exposure prophylaxis)	X	X	X		
Rotavirus (oral)	X	Х	X		
Tuberculosis	X	X	X		
Typhoid fever	X	X	Х		
Typhoid fever and Hepatitis A combination	X	X	X		
Yellow fever	X	X	X		

The information in this document is current as of the publication date. This information serves as a guide only and should not be employed as a substitute or replacement for reviewing regulations or as a clinical decision-making tool. Published 13 March 2023